

TEST BANK


# Pharmacotherapeutics

## for Advanced Practice Nurse Prescribers

Teri Moser Woo, Marylou V. Robinson

5th Edition

Woo • Robinson



# Table of Contents

## **Unit I. The Foundation**

- Chapter 1: The Role of the Advanced Practice Nurse as Prescriber
- Chapter 2: Review of Basic Principles of Pharmacology
- Chapter 3: Rational Drug Selection
- Chapter 4: Legal and Professional Issues in Prescribing
- Chapter 5: Adverse Drug Reactions
- Chapter 6: An Introduction to Pharmacogenomics
- Chapter 7: Nutrition and Nutraceuticals
- Chapter 8: Herbal Therapies and Cannabis
- Chapter 9: Information Technology and Pharmacotherapeutics
- Chapter 10: Pharmacoeconomics

## **Unit II. Pharmacotherapeutics with Single Drugs**

- Chapter 11: Drugs Affecting the Autonomic Nervous System
- Chapter 12: Drugs Affecting the Central Nervous System
- Chapter 13: Drugs Affecting the Cardiovascular and Renal Systems
- Chapter 14: Drugs Affecting the Respiratory System
- Chapter 15: Drugs Affecting the Hematological System
- Chapter 16: Drugs Affecting the Immune System
- Chapter 17: Drugs Affecting the Gastrointestinal System
- Chapter 18: Drugs Affecting the Endocrine System: Pancreatic Hormones and Antidiabetic Drugs
- Chapter 19: Drugs Affecting the Endocrine System: Pituitary, Thyroid, and Adrenal Drugs
- Chapter 20: Drugs Affecting the Reproductive System
- Chapter 21: Drugs Affecting the Bones and Joints
- Chapter 22: Drugs Affecting the Integumentary System
- Chapter 23: Drugs Used to Treat Bacterial Infections
- Chapter 24: Drugs Used to Treat Viral, Fungal, and Protozoal Infections
- Chapter 25: Drugs Used to Treat Inflammatory Processes
- Chapter 26: Drugs Used to Treat Eye and Ear Disorders

## **Unit III. Pharmacotherapeutics with Multiple Drugs**

- Chapter 27: Alcohol and Drug Addiction
- Chapter 28: Anemia
- Chapter 29: Angina
- Chapter 30: Anxiety and Depression
- Chapter 31: Attention Deficit-Hyperactivity Disorder
- Chapter 32: Asthma and Chronic Obstructive Pulmonary Disease
- Chapter 33 : Contraception
- Chapter 34 : Dermatological Conditions
- Chapter 35 : Diabetes Mellitus
- Chapter 36: Gastroesophageal Reflux and Peptic Ulcer Disease
- Chapter 37: Headaches
- Chapter 38: Heart Failure
- Chapter 39: Human Immunodeficiency Virus Disease and Acquired Immunodeficiency Syndrome
- Chapter 40: Hormone Replacement Therapy
- Chapter 41: Hyperlipidemia
- Chapter 42: Hypertension
- Chapter 43: Hyperthyroidism and Hypothyroidism
- Chapter 44: Pain Management: Acute and Chronic Pain
- Chapter 45: Pneumonia
- Chapter 46: Tobacco Cessation
- Chapter 47: Sexually Transmitted Infections and Vaginitis
- Chapter 48: Tuberculosis
- Chapter 49: Upper Respiratory Infection, Pharyngitis, Sinusitis, Otitis Media, and Otitis Externa
- Chapter 50: Urinary Tract Infections

## **Unit IV. Special Drug Treatment Considerations**

- Chapter 51: Women as Patients
- Chapter 52: Men as Patients
- Chapter 53: Pediatric Patients
- Chapter 54: Transgendered Clients as Patients
- Chapter 55: Geriatric Patients

## Chapter 1. The Role of the Nurse Practitioner

### Multiple Choice

*Identify the choice that best completes the statement or answers the question.*

1. Nurse practitioner prescriptive authority is regulated by:

1. The National Council of State Boards of Nursing
2. The U.S. Drug Enforcement Administration
3. The State Board of Nursing for each state
4. The State Board of Pharmacy

2. The benefits to the patient of having an Advanced Practice Registered Nurse (APRN) prescriber include:

1. Nurses know more about Pharmacology than other prescribers because they take it both in their basic nursing program and in their APRN program.
2. Nurses care for the patient from a holistic approach and include the patient in decision making regarding their care.
3. APRNs are less likely to prescribe narcotics and other controlled substances.
4. APRNs are able to prescribe independently in all states, whereas a physician's assistant needs to have a physician supervising their practice.

3. Clinical judgment in prescribing includes:

1. Factoring in the cost to the patient of the medication prescribed
2. Always prescribing the newest medication available for the disease process
3. Handing out drug samples to poor patients
4. Prescribing all generic medications to cut costs

4. Criteria for choosing an effective drug for a disorder include:

1. Asking the patient what drug they think would work best for them
2. Consulting nationally recognized guidelines for disease management
3. Prescribing medications that are available as samples before writing a prescription
4. Following U.S. Drug Enforcement Administration guidelines for prescribing

5. Nurse practitioner practice may thrive under health-care reform because of:

1. The demonstrated ability of nurse practitioners to control costs and improve patient outcomes
2. The fact that nurse practitioners will be able to practice independently
3. The fact that nurse practitioners will have full reimbursement under health-care reform
4. The ability to shift accountability for Medicaid to the state level

## Chapter 1. The Role of the Nurse Practitioner

### Answer Section

#### MULTIPLE CHOICE

1. ANS: 3	PTS: 1
2. ANS: 2	PTS: 1
3. ANS: 1	PTS: 1
4. ANS: 2	PTS: 1
5. ANS: 1	PTS: 1

## Chapter 2. Review of Basic Principles of Pharmacology

### Multiple Choice

*Identify the choice that best completes the statement or answers the question.*

- \_\_\_\_ 1. A patient's nutritional intake and laboratory results reflect hypoalbuminemia. This is critical to prescribing because:
  1. Distribution of drugs to target tissue may be affected.
  2. The solubility of the drug will not match the site of absorption.
  3. There will be less free drug available to generate an effect.
  4. Drugs bound to albumin are readily excreted by the kidneys.
- \_\_\_\_ 2. Drugs that have a significant first-pass effect:
  1. Must be given by the enteral (oral) route only
  2. Bypass the hepatic circulation
  3. Are rapidly metabolized by the liver and may have little if any desired action
  4. Are converted by the liver to more active and fat-soluble forms
- \_\_\_\_ 3. The route of excretion of a volatile drug will likely be the:
  1. Kidneys
  2. Lungs
  3. Bile and feces
  4. Skin
- \_\_\_\_ 4. Medroxyprogesterone (Depo Provera) is prescribed intramuscularly (IM) to create a storage reservoir of the drug. Storage reservoirs:
  1. Assure that the drug will reach its intended target tissue
  2. Are the reason for giving loading doses
  3. Increase the length of time a drug is available and active
  4. Are most common in collagen tissues
- \_\_\_\_ 5. The NP chooses to give cephalexin every 8 hours based on knowledge of the drug's:
  1. Propensity to go to the target receptor
  2. Biological half-life
  3. Pharmacodynamics
  4. Safety and side effects

6. Azithromycin dosing requires that the first day's dosage be twice those of the other 4 days of the prescription. This is considered a loading dose. A loading dose:

1. Rapidly achieves drug levels in the therapeutic range
2. Requires four- to five-half-lives to attain
3. Is influenced by renal function
4. Is directly related to the drug circulating to the target tissues

7. The point in time on the drug concentration curve that indicates the first sign of a therapeutic effect is the:

1. Minimum adverse effect level
2. Peak of action
3. Onset of action
4. Therapeutic range

8. Phenytoin requires that a trough level be drawn. Peak and trough levels are done:

1. When the drug has a wide therapeutic range
2. When the drug will be administered for a short time only
3. When there is a high correlation between the dose and saturation of receptor sites
4. To determine if a drug is in the therapeutic range

9. A laboratory result indicates that the peak level for a drug is above the minimum toxic concentration. This means that the:

1. Concentration will produce therapeutic effects
2. Concentration will produce an adverse response
3. Time between doses must be shortened
4. Duration of action of the drug is too long

10. Drugs that are receptor agonists may demonstrate what property?

1. Irreversible binding to the drug receptor site
2. Upregulation with chronic use
3. Desensitization or downregulation with continuous use
4. Inverse relationship between drug concentration and drug action

11. Drugs that are receptor antagonists, such as beta blockers, may cause:

1. Downregulation of the drug receptor
2. An exaggerated response if abruptly discontinued
3. Partial blockade of the effects of agonist drugs
4. An exaggerated response to competitive drug agonists

12. Factors that affect gastric drug absorption include:

1. Liver enzyme activity
2. Protein-binding properties of the drug molecule
3. Lipid solubility of the drug
4. Ability to chew and swallow

13. Drugs administered via IV:

1. Need to be lipid soluble in order to be easily absorbed
2. Begin distribution into the body immediately
3. Are easily absorbed if they are nonionized
4. May use pinocytosis to be absorbed

14. When a medication is added to a regimen for a synergistic effect, the combined effect of the drugs is:
  1. The sum of the effects of each drug individually
  2. Greater than the sum of the effects of each drug individually
  3. Less than the effect of each drug individually
  4. Not predictable, as it varies with each individual
  
15. Which of the following statements about bioavailability is true?
  1. Bioavailability issues are especially important for drugs with narrow therapeutic ranges or sustained-release mechanisms.

2. All brands of a drug have the same bioavailability.
3. Drugs that are administered more than once a day have greater bioavailability than drugs given once daily.
4. Combining an active drug with an inert substance does not affect bioavailability.

16. Which of the following statements about the major distribution barriers (blood-brain or fetal-placental) is true?

1. Water soluble and ionized drugs cross these barriers rapidly.
2. The blood-brain barrier slows the entry of many drugs into and from brain cells.
3. The fetal-placental barrier protects the fetus from drugs taken by the mother.
4. Lipid-soluble drugs do not pass these barriers and are safe for pregnant women.

17. Drugs are metabolized mainly by the liver via phase I or phase II reactions. The purpose of both of these types of reactions is to:

1. Inactivate prodrugs before they can be activated by target tissues
2. Change the drugs so they can cross plasma membranes
3. Change drug molecules to a form that an excretory organ can excrete
4. Make these drugs more ionized and polar to facilitate excretion

18. Once they have been metabolized by the liver, the metabolites may be:

1. More active than the parent drug
2. Less active than the parent drug
3. Totally “deactivated” so they are excreted without any effect
4. All of the above

19. All drugs continue to act in the body until they are changed or excreted. The ability of the body to excrete drugs via the renal system would be increased by:

1. Reduced circulation and perfusion of the kidney
2. Chronic renal disease
3. Competition for a transport site by another drug
4. Unbinding a nonvolatile drug from plasma proteins

20. Steady state is:

1. The point on the drug concentration curve when absorption exceeds excretion
2. When the amount of drug in the body remains constant
3. When the amount of drug in the body stays below the minimum toxic concentration
4. All of the above

21. Two different pain medications are given together for pain relief. The drug—drug interaction is:

1. Synergistic
2. Antagonistic
3. Potentiative
4. Additive

22. Actions taken to reduce drug—drug interaction problems include all of the following EXCEPT:

1. Reducing the dosage of one of the drugs
2. Scheduling their administration at different times
3. Prescribing a third drug to counteract the adverse reaction of the combination

4. Reducing the dosage of both drugs

23. Phase I oxidative-reductive processes of drug metabolism require certain nutritional elements. Which of the following would reduce or inhibit this process?

1. Protein malnutrition
2. Iron-deficiency anemia
3. Both 1 and 2
4. Neither 1 nor 2

24. The time required for the amount of drug in the body to decrease by 50% is called:

1. Steady state
2. Half-life
3. Phase II metabolism
4. Reduced bioavailability time

25. An agonist activates a receptor and stimulates a response. When given frequently over time, the body may:

1. Upregulate the total number of receptors
2. Block the receptor with a partial agonist
3. Alter the drug's metabolism
4. Downregulate the numbers of that specific receptor

26. Drug antagonism is best defined as an effect of a drug that:

1. Leads to major physiological and psychological dependence
2. Is modified by the concurrent administration of another drug
3. Cannot be metabolized before another dose is administered
4. Leads to a decreased physiological response when combined with another drug

27. Instructions to a client regarding self-administration of oral enteric-coated tablets should include which of the following statements?

1. "Avoid any other oral medicines while taking this drug."
2. "If swallowing this tablet is difficult, dissolve it in 3 ounces of orange juice."
3. "The tablet may be crushed if you have any difficulty taking it."
4. "To achieve best effect, take the tablet with at least 8 ounces of fluid."

28. The major reason for not crushing a sustained-release capsule is that, if crushed, the coated beads of the drugs could possibly result in:

1. Disintegration
2. Toxicity
3. Malabsorption
4. Deterioration

29. Which of the following substances is the most likely to be absorbed in the intestines rather than in the stomach?

1. Sodium bicarbonate
2. Ascorbic acid
3. Salicylic acid
4. Glucose

\_\_\_\_\_ 30. Which of the following variables is a factor in drug absorption?

1. The smaller the surface area for absorption, the more rapidly the drug is absorbed.
2. A rich blood supply to the area of absorption leads to better absorption.
3. The less soluble the drug, the more easily it is absorbed.
4. Ionized drugs are easily absorbed across the cell membrane.

\_\_\_\_\_ 31. An advantage of prescribing a sublingual medication is that the medication is:

1. Absorbed rapidly
2. Excreted rapidly
3. Metabolized minimally
4. Distributed equally

\_\_\_\_\_ 32. Drugs that use CYP 3A4 isoenzymes for metabolism may:

1. Induce the metabolism of another drug
2. Inhibit the metabolism of another drug
3. Both 1 and 2
4. Neither 1 nor 2

\_\_\_\_\_ 33. Therapeutic drug levels are drawn when a drug reaches steady state. Drugs reach steady state:

1. After the second dose
2. After four to five half-lives
3. When the patient feels the full effect of the drug
4. One hour after IV administration

\_\_\_\_\_ 34. Upregulation or hypersensitization may lead to:

1. Increased response to a drug
2. Decreased response to a drug
3. An exaggerated response if the drug is withdrawn
4. Refractoriness or complete lack of response

## Chapter 2. Review of Basic Principles of Pharmacology

### Answer Section

#### MULTIPLE CHOICE

1. ANS 1	PTS: 1
:	
2. ANS 3	PTS: 1
:	
3. ANS 2	PTS: 1
:	
4. ANS 3	PTS: 1
:	
5. ANS 2	PTS: 1
:	
6. ANS 1	PTS: 1
:	
7. ANS 3	PTS: 1
:	
8. ANS 4	PTS: 1
:	
9. ANS 2	PTS: 1
:	
10. ANS 3	PTS: 1
:	
11. ANS 2	PTS: 1
:	
12. ANS 3	PTS: 1
:	
13. ANS 2	PTS: 1
:	
14. ANS 2	PTS: 1
:	
15. ANS 1	PTS: 1
:	
16. ANS 2	PTS: 1
:	
17. ANS 3	PTS: 1
:	
18. ANS 4	PTS: 1
:	
19. ANS 4	PTS: 1
:	
20. ANS 2	PTS: 1
:	
21. ANS 4	PTS: 1
:	
22. ANS 3	PTS: 1
:	
23. ANS 4	PTS: 1
:	