

TEST BANK

Berne & Levy Physiology

Bruce M. Koeppen, Bruce A. Stanton, Julianne M. Hall, Agnieszka Swiatecka-Urban

8th Edition




Table of Contents

Section 1: Cellular Physiology

- 1 Principles of Cell and Membrane Function
- 2 Homeostasis: Volume and Composition of Body Fluid Compartments
- 3 Signal Transduction, Membrane Receptors, Second Messengers, and Regulation of Gene Expression

Section 2: Neurophysiology

- 4 The Nervous System: Introduction to Cells and Systems
- 5 Generation and Conduction of Action Potentials
- 6 Synaptic Transmission
- 7 The Somatosensory System
- 8 The Special Senses
- 9 Organization of Motor Function
- 10 Integrative Functions of the Nervous System
- 11 The Autonomic Nervous System and Its Central Control

Section 3: Muscle Physiology

- 12 Skeletal Muscle Physiology
- 13 Cardiac Muscle
- 14 Smooth Muscle

Section 4: Cardiovascular Physiology

- 15 Overview of Circulation
- 16 Elements of Cardiac Function
- 17 Properties of the Vasculature
- 18 Regulation of the Heart and Vasculature
- 19 Integrated Control of the Cardiovascular System

Section 5: Respiratory Physiology

- 20 Introduction to the Respiratory System
- 21 Static Lung and Chest Wall Mechanics
- 22 Dynamic Lung and Chest Wall Mechanics
- 23 Ventilation, Perfusion, and Ventilation/Perfusion Relationships
- 24 Oxygen and Carbon Dioxide Transport
- 25 Control of Respiration
- 26 Host Defense and Metabolism in the Lung

Section 6: Gastrointestinal Physiology

- 27 Functional Anatomy and General Principles of Regulation in the Gastrointestinal Tract
- 28 The Cephalic, Oral, and Esophageal Phases of the Integrated Response to a Meal
- 29 The Gastric Phase of the Integrated Response to a Meal
- 30 The Small Intestinal Phase of the Integrated Response to a Meal
- 31 The Colonic Phase of the Integrated Response to a Meal
- 32 Transport and Metabolic Functions of the Liver

Section 7: Renal Physiology

- 33 Elements of Renal Function
- 34 Solute and Water Transport Along the Nephron: Tubular Function
- 35 Control of Body Fluid Osmolality and Volume
- 36 Potassium, Calcium, and Phosphate Homeostasis
- 37 Role of the Kidneys in the Regulation of Acid-Base Balance

Section 8: Endocrine Physiology

- 38 Introduction to the Endocrine System
- 39 Hormonal Regulation of Energy Metabolism
- 40 Hormonal Regulation of Calcium and Phosphate Metabolism
- 41 The Hypothalamus and Pituitary Gland
- 42 The Thyroid Gland
- 43 The Adrenal Gland
- 44 The Male and Female Reproductive Systems


Chapter 01: Principles of Cell Function
Koeppen: Berne and Levy Physiology, 8th Edition

MULTIPLE CHOICE

1. The subcellular structure that degrades proteins is called the:
 - a. Tight junction
 - b. Mitochondria
 - c. Lysosome
 - d. Plasma membrane
 - e. Ribosome

ANS: C PTS: 1

2. An experiment is done to measure the uptake of an amino acid into a cell. The following data are obtained:




If Na^+ is removed from the extracellular bathing solution, or if a drug is added that prevents the cell from making adenosine triphosphate (ATP), the uptake of amino acid into the cell is markedly reduced. According to this information, which of the following mechanisms is probably responsible for the transport of the amino acid into the cell?

- a. Passive diffusion through the lipid bilayer
- b. Uniporter
- c. Transport ATPase
- d. Na^+ symporter
- e. Na^+ antiporter

ANS: D PTS: 1

3. A membrane permeable by only Na^+ separates two compartments containing Na_2SO_4 , shown as follows:



Electrodes are placed in both compartments, and a voltage is applied (that of compartment A is held at 0 mV). What voltage applied to compartment B would result in *no* net movement of Na^+ across the membrane separating the compartments?

- a. -60 mV
- b. -30 mV
- c. 0 mV
- d. +30 mV
- e. +60 mV

ANS: E PTS: 1

4. The resting membrane potential of a cell is -85 mV. The intracellular and extracellular concentrations of several ions are indicated in the following table, as is the calculated Nernst equilibrium potential (E_i) for each of these ions:

Ion	Concentration Inside Cell	Concentration Outside Cell	E_i
Na^+	12 mEq/L	145 mEq/L	66 mV
K^+	150 mEq/L	4 mEq/L	-96 mV
Cl^-	30 mEq/L	105 mEq/L	-33 mV
Ca^{++}	0.0001 mmol/dL	1 mmol/dL	122 mV

The membrane has channels for Na^+ , K^+ , Cl^- , and Ca^{++} . The conductance of the membrane is the greatest for which ion?

- a. Na^+
- b. K^+
- c. Cl^-
- d. Ca^{++}

ANS: B PTS: 1

5. A cell contains the following membrane transporters:

Na^+ channel
 K^+ channel
 Na^+, K^+ -ATPase

The resting membrane voltage of the cell is -80 mV, and the intracellular and extracellular ion concentrations are as follows:

Ion	Intracellular Concentration	Extracellular Concentration
Na^+	10 mEq/L	145 mEq/L
K^+	120 mEq/L	4 mEq/L

The cell is treated with a drug to inhibit the Na^+, K^+ -ATPase. What would be the effect of this drug on the following parameters?


	Intracellular $[\text{Na}^+]$	Intracellular	Cell Volume	Membrane Voltage
--	-------------------------------	---------------	-------------	------------------

		[K ⁺]		
A.	Decrease	Decrease	Decrease	Depolarize
B.	Increase	Decrease	Increase	Depolarize
C.	Increase	Increase	Increase	No change
D.	Decrease	Increase	Decrease	Hyperpolarize
E.	Increase	Increase	Decrease	Hyperpolarize

- A.
- B.
- C.
- D.
- E.

ANS: B PTS: 1

6. Which of the labeled proteins (shaded) is attached to the membrane by a glycosylphosphatidylinositol (GPI) anchor?



- A
- B
- C
- D
- E

ANS: E PTS: 1

7. A cell has channels for Na⁺, K⁺, and Cl⁻ in its plasma membrane. The resting membrane potential is -60 mV (cell interior negative). The intracellular and extracellular concentrations for these ions, as well as the calculated Nernst potentials, are listed as follows:

Ion	Intracellular Concentration	Extracellular Concentration	E _i
Na ⁺	14 mEq/L	140 mEq/L	60 mV
K ⁺	150 mEq/L	5 mEq/L	-89 mV
Cl ⁻	10 mEq/L	100 mEq/L	-60 mV

A drug is applied to the cell that increases the permeability of the cell by Cl⁻ (i.e., it opens Cl⁻ channels). What effect will this drug have on the net movement of Cl⁻ across the plasma membrane?

- Net Cl⁻ movement out of the cell will be increased.